



MONSON
VISION

CATARACT POST-OP INSTRUCTIONS


POST-OPERATIVE MEDICATIONS

PRED-MOXI-BROM SUSPENSION



Shake well before using.
Place 1 drop in surgical eye 4 times daily, for 2 weeks, followed by 2 times daily for the next 2 weeks.

SURGERY EYE DROP SCHEDULE





MONSON
VISION

435.787.7200

CATARACT POST-OP INSTRUCTIONS

WHAT TO EXPECT AFTER SURGERY



FIRST DAY

- Start your post-op drops immediately.
- A medication was placed in your eye during surgery to reduce your dilated pupil. This medication may cause a temporary headache, and your pupil may appear smaller for a day or two. This is normal.
- Your vision may be blurry on your treatment day. It is normal to experience light sensitivity, halos, glare, dryness, fluctuation in vision, and the sensation that something is in your eye.



FIRST WEEK

- Blurry vision will slowly improve over the first week.
- You may notice some red spots on the white part of your eye. These will resolve as your eye heals.
- You may experience tenderness following the procedure. This is normal, and will improve the first week.
- Wear the clear eye shield taped over the eyebrow while sleeping for 7 days.
- Avoid lifting over 25 pounds.
- Avoid rubbing eyes.
- Avoid strenuous activity.
- Avoid getting water in your eyes for 1 week.



WHEN TO CALL YOUR SURGEON

- Mild discomfort and aching is normal. If you have severe pain that does not improve with 2 extra strength Tylenol, please call the office immediately.
- If you have questions or concerns about eye drops.
- If you experience flashes of lights, new floaters or loss of peripheral vision please call immediately.
- It is our policy at Monson Vision, that a surgical patient who calls with any concern will be given the option of coming into the office immediately to be seen.



POST-OP APPOINTMENTS

1st eye: Day 1 _____ Week 1 _____ Dr. _____
 2nd eye: Day 1 _____ Week 1 _____ Dr. _____